

Sections Affected in the Finding Aids section of this volume.

EDITORIAL NOTE: At 62 FR 29014, May 29, 1997, in § 558.550, paragraph (b) was redesignated as paragraph (d). At 62 FR 60781, Nov. 13, 1997, paragraph (b)(1)(vii)(c) was amended by removing "No. 000004" and adding in its place "Nos. 000004 and 046573" and at 62 FR 66985, Dec. 23, 1997, paragraph (b)(1)(ix)(c) was amended by removing "No. 000004" and adding in its place "Nos. 000004 and 046573". Since § 558.550(b) does not exist, these amendments could not be incorporated.

§ 558.555 Semduramicin.

(a) *Approvals.* Type A medicated article containing 5.13 percent semduramicin sodium (equivalent to 50 grams semduramicin per kilogram or 22.7 grams per pound) to 000069 in § 510.600(c) of this chapter.

(b) *Conditions of use.* (1) Broilers:

(i) *Amount.* Semduramicin: 25 parts per million.

(ii) *Indications for use.* For the prevention of coccidiosis caused by *Eimeria tenella*, *E. acervulina*, *E. maxima*, *E. brunetti*, *E. necatrix*, and *E. mivati/E. mitis*.

(iii) *Limitations.* Do not feed to laying hens.

(2) *Amount.* Semduramicin 22.7 grams with bacitracin methylene disalicylate 10 to 50 grams and roxarsone 45.4 grams per ton.

(i) *Indications for use.* For the prevention of coccidiosis caused by *Eimeria acervulina*, *E. brunetti*, *E. maxima*, *E. mivati/E. mitis*, *E. necatrix*, and *E. tenella*, including some field strains of *E. tenella* that are more susceptible to semduramicin combined with roxarsone than semduramicin alone, and for improved feed efficiency.

(ii) *Limitations.* Feed continuously as sole ration. Use feed within 2 weeks of production. Withdraw 5 days before slaughter. Do not feed to laying hens. Use as sole source of organic arsenic. Poultry should have access to drinking water at all times. Drug overdose or lack of water intake may result in leg weakness or paralysis. Semduramicin as provided by 000069 in § 510.600(c) of this chapter, bacitracin methylene disalicylate and roxarsone as provided by 046573.

(3) *Amount.* Semduramicin 22.7 grams with bacitracin methylene disalicylate 10 to 50 grams per ton.

(i) *Indications for use.* For the prevention of coccidiosis caused by *Eimeria acervulina*, *E. brunetti*, *E. maxima*, *E. mivati/E. mitis*, *E. necatrix*, and *E. tenella*, and for improved feed efficiency in broiler chickens.

(ii) *Limitations.* Feed continuously as sole ration. Use feed within 2 weeks of production. Do not feed to laying hens. Semduramicin as provided by 000069, bacitracin methylene disalicylate as provided by 046573 in § 510.600(c) of this chapter.

(4) *Amount.* Semduramicin 22.7 grams with roxarsone 45.4 grams per ton.

(i) *Indications for use.* For the prevention of coccidiosis caused by *Eimeria acervulina*, *E. brunetti*, *E. maxima*, *E. mivati/E. mitis*, *E. necatrix*, and *E. tenella*, including some field strains of *E. tenella* that are more susceptible to semduramicin combined with roxarsone than semduramicin alone.

(ii) *Limitations.* Feed continuously as sole ration. Withdraw 5 days before slaughter. For broiler chickens only. Do not feed to laying hens. Use as sole source of organic arsenic. Roxarsone as provided by 046573, semduramicin as provided by 000069 in § 510.600(c) of this chapter.

[59 FR 17477, Apr. 13, 1994, as amended at 60 FR 57928, Nov. 24, 1995; 61 FR 29481, June 11, 1996; 61 FR 43451, Aug. 23, 1996; 61 FR 66584, Dec. 18, 1996; 62 FR 66985, Dec. 23, 1997]

§ 558.575 Sulfadimethoxine, ormetoprim.

(a) *Approvals.* Type A medicated articles to sponsors as identified in § 510.600(c) of this chapter for uses as in paragraph (c) of this section as follows:

(1) 25 percent sulfadimethoxine and 15 percent ormetoprim to 000004 for use for poultry as in paragraphs (c)(1), (2), (3), and (4) of this section.

(2) 25 percent sulfadimethoxine and 5 percent ormetoprim to 000004 for use for fish as in paragraphs (c)(5) and (6) of this section.

(b) *Related tolerances.* See §§ 556.490 and 556.640 of this chapter.

(c) *Conditions of use.* It is used in feeds for animals as follows:

(1) *Broiler chickens*—(i) *Amount per ton.* Sulfadimethoxine, 113.5 grams (0.0125 percent) plus ormetoprim, 68.1 grams (0.0075 percent).